

National Buruli Ulcer Control Programme-Ghana
Summary Regional Reported Cases of Buruli Ulcer: Jan-Dec 2009

Region	Total New Cases	Total Population	Prevalence/100000	Sex		Age Group				Limitation of joint movement		Referred BY						Clinical Form				Category of Lesion			Location of Lesion							Proportion Specimen
				Female	Male	0-5 yrs	6-15 yrs	16-49 yrs	Age >50yrs	No	Yes	CV	FP	HW	OT	S	ST	Nodule	Oedema	Plaque	Ulcer	Cat I	Cat II	Cat III	Abdomen	Back	Buttocks & perineum	Nead & Neck	Lower Limb(s)	Thorax	Upper Limb(s)	
Ashanti	428	4,839,100	8.84	231	197	49	139	132	103	335	91	83	29	103	46	154	5	62	45	43	249	145	143	135	12	3	4	10	277	6	115	352/416
Brong Ahafo	66	2,257,304	2.92	35	31	0	21	31	14	57	9	9	0	18	2	13	1	13	6	2	28	36	9	0	1	0	1	0	45	0	18	18/51
Central	162	1,864,104	8.69	90	72	15	28	63	56	62	100	8	89	34	0	31	0	10	55	5	90	7	50	105	1	1	0	2	131	0	25	77/157
Eastern	65	2,297,565	2.83	33	32	2	17	20	26	56	7	10	0	21	5	22	0	2	13	1	47	23	27	12	0	2	1	3	47	1	9	38/54
Greater Accra	113	4,358,263	2.59	66	47	11	40	49	13	81	32	20	2	26	11	52	2	17	4	4	81	39	24	50	2	2	1	3	66	7	31	73/113
Western	17	2,558,113	0.66	10	7	0	1	13	3	14	2	1	1	6	1	6	0	1	0	0	13	3	6	3	0	0	0	1	10	0	2	3/7
National Total	851	23,951,519	3.55	465	386	77	246	308	215	605	241	131	121	208	65	278	8	105	123	55	508	253	259	305	16	8	7	19	576	14	200	561/798